arXiv:1611.01413v1 [math.GM] 26 Oct 2016

A geometrization of quadratic multi-time Lagrangians

Mircea Neagu

Abstract

The aim of this paper is to construct a Riemann-Lagrange geometry (in the sense of d-linear connection, d-torsions and d-curvatures) for a quadratic multi-time Lagrangian.

Mathematics Subject Classifications (2010): 58B20, 53C60, 53C80.
Key words and phrases: 1-jet spaces, quadratic multi-time Lagrangian, nonlinear connection, Cartan linear d-connection, d-torsions and d-curvatures.

1 Introduction

It is important to note that the quadratic multi-time Lagrangians dominates most physical domains. We can remind only the theory of elasticity [7], the dynamics of ideal fluids, the magnetohydrodynamics [3], [4] or the theory of bosonic strings [2]. This fact emphasizes the importance of the geometrization of quadratic multi-time Lagrangians. In conclusion, a corresponding Riemann-Lagrange geometry on 1-jet spaces is required.

2 The generalized multi-time Lagrange space of a quadratic Lagrangian

Let $U_{(i)}^{(\alpha)}(t^\gamma, x^k)$ be a d-tensor on the 1-jet space $J^1(T, M)$ and let $F : T \times M \to \mathbb{R}$ be a smooth function. Let us consider the quadratic multi-time Lagrangian function $L : J^1(T, M) \to \mathbb{R}$, defined by

$$L = G_{(i)(j)}^{(\alpha)(\beta)}(t^\gamma, x^k) x_\alpha^i x_\beta^j + U_{(i)}^{(\alpha)}(t^\gamma, x^k) x_\alpha^i + F(t^\gamma, x^k), \tag{1}$$

whose fundamental vertical metrical d-tensor

$$G_{(i)(j)}^{(\alpha)(\beta)}(t^\gamma, x^k) = \frac{1}{2} \frac{\partial^2 L}{\partial x_\alpha^i \partial x_\beta^j}$$

is symmetric, of rank $n = \dim M$ and has a constant signature with respect to the indices i and j. Using a semi-Riemannian metric $h = (h_{\alpha\beta}(t^\gamma))_{\alpha,\beta=\overline{1,p}}$ on

T and considering the canonical Kronecker h-regular vertical metrical d-tensor attached to the Lagrangian function (1), given by

$$\mathcal{G}_{(i)(j)}^{(\alpha)(\beta)}(t^\gamma, x^k) = \frac{1}{p} h^{\alpha\beta}(t^\gamma) h_{\mu\nu}(t^\gamma) G_{(i)(j)}^{(\mu)(\nu)}(t^\gamma, x^k),$$

where $p = \dim T$, then the pair

$$\mathcal{QLG}ML_p^n = \left(J^1(T, M), \mathcal{G}_{(i)(j)}^{(\alpha)(\beta)}(t^\gamma, x^k) = h^{\alpha\beta}(t^\gamma) g_{ij}(t^\gamma, x^k) \right)$$

is a generalized multi-time Lagrange space whose spatial metrical d-tensor is given by the formula

$$g_{ij}(t^\gamma, x^k) = \frac{1}{p} h_{\mu\nu}(t^\gamma) G_{(i)(j)}^{(\mu)(\nu)}(t^\gamma, x^k). \tag{2}$$

Definition 1 *The space $\mathcal{QLG}ML_p^n$ is called the* ***canonical generalized multi-time Lagrange space associated with the quadratic Lagrangian function (1)****.*

In order to develop the Riemann-Lagrange geometry of the space $\mathcal{QLG}ML_p^n$ (in the sense of d-linear connection, d-torsions and d-curvatures) we need a nonlinear connection $\Gamma = \left(M_{(\alpha)\beta}^{(i)}, N_{(\alpha)j}^{(i)} \right)$ on $J^1(T, M)$. The fundamental vertical metrical d-tensor $\mathcal{G}_{(i)(j)}^{(\alpha)(\beta)}(t^\gamma, x^k) = h^{\alpha\beta}(t^\gamma) g_{ij}(t^\gamma, x^k)$, where g_{ij} is given by (2), produces the following natural nonlinear connection (see [6], pp. 88):

$$M_{(\alpha)\beta}^{(i)} = -H_{\alpha\beta}^{\gamma} x_\gamma^i, \quad N_{(\alpha)j}^{(i)} = \Gamma_{jm}^i x_\alpha^m + \frac{g^{im}}{2} \frac{\partial g_{jm}}{\partial t^\alpha},$$

where $H_{\alpha\beta}^{\gamma}$ are the Christoffel symbols of the temporal semi-Riemannian metric $h_{\alpha\beta}$, and

$$\Gamma_{jk}^i(t^\mu, x^m) = \frac{g^{ir}}{2} \left(\frac{\partial g_{jr}}{\partial x^k} + \frac{\partial g_{kr}}{\partial x^j} - \frac{\partial g_{jk}}{\partial x^r} \right)$$

are the generalized Christoffel symbols of the spatial metric $g_{ij}(t^\gamma, x^k)$. Let

$$\left\{ \frac{\delta}{\delta t^\alpha}, \frac{\delta}{\delta x^i}, \frac{\partial}{\partial x_\alpha^i} \right\} \subset \mathcal{X}\left(J^1(T, M) \right) \text{ and } \left\{ dt^\alpha, dx^i, \delta x_\alpha^i \right\} \subset \mathcal{X}^*\left(J^1(T, M) \right)$$

be the adapted bases of the nonlinear connection Γ, where

$$\frac{\delta}{\delta t^\alpha} = \frac{\partial}{\partial t^\alpha} - M_{(\beta)\alpha}^{(j)} \frac{\partial}{\partial x_\beta^j}, \quad \frac{\delta}{\delta x^i} = \frac{\partial}{\partial x^i} - N_{(\beta)i}^{(j)} \frac{\partial}{\partial x_\beta^j},$$

$$\delta x_\alpha^i = dx_\alpha^i + M_{(\alpha)\beta}^{(i)} dt^\beta + N_{(\alpha)j}^{(i)} dx^j.$$

Remark 2 *In what follows, all geometrical objects that we construct will be described in local adapted components with respect to previous adapted bases.*

3 The Riemann-Lagrange geometry of the space $\mathcal{QLG}ML_p^n$

Following the geometrical ideas from the works [6], [1], [5], we can enunciate the main results of the Riemann-Lagrange geometry of the generalized multi-time Lagrange space $\mathcal{QLG}ML_p^n$.

Theorem 3 (the Cartan linear connection) *The canonical Cartan linear connection of the space $\mathcal{QLG}ML_p^n$ is given by the adapted components*

$$C\Gamma = \left(H_{\alpha\beta}^{\gamma}, G_{j\gamma}^{k}, L_{jk}^{i} = \Gamma_{jk}^{i}, C_{j(k)}^{i(\gamma)} = 0\right),$$

where $G_{j\gamma}^{k} = \dfrac{g^{km}}{2}\dfrac{\partial g_{mj}}{\partial t^{\gamma}}$.

Proof. The formulas which generally describe the adapted coefficients of the Cartan canonical connection are given by

$$G_{j\gamma}^{k} = \frac{g^{km}}{2}\frac{\delta g_{mj}}{\delta t^{\gamma}} = \frac{g^{km}}{2}\frac{\partial g_{mj}}{\partial t^{\gamma}}, \quad L_{jk}^{i} = \frac{g^{im}}{2}\left(\frac{\delta g_{jm}}{\delta x^{k}} + \frac{\delta g_{km}}{\delta x^{j}} - \frac{\delta g_{jk}}{\delta x^{m}}\right) = \Gamma_{jk}^{i},$$

$$C_{j(k)}^{i(\gamma)} = \frac{g^{im}}{2}\left(\frac{\partial g_{jm}}{\partial x_{\gamma}^{k}} + \frac{\partial g_{km}}{\partial x_{\gamma}^{j}} - \frac{\partial g_{jk}}{\partial x_{\gamma}^{m}}\right) = 0.$$

■

Remark 4 *The generalized Cartan canonical connection $C\Gamma$ of the generalized multi-time Lagrange space $\mathcal{QLG}ML_p^n$ satisfies the metrical conditions*

$$h_{\alpha\beta/\gamma} = h_{\alpha\beta|k} = h_{\alpha\beta}|_{(k)}^{(\gamma)} = 0, \qquad g_{ij/\gamma} = g_{ij|k} = g_{ij}|_{(k)}^{(\gamma)} = 0,$$

where "$_{/\gamma}$", "$_{|k}$" and "$|_{(k)}^{(\gamma)}$" are the local covariant derivatives produced by the Cartan connection $C\Gamma$.

Theorem 5 *The generalized multi-time Lagrange space $\mathcal{QLG}ML_p^n$ is characterized by the following adapted torsion d-tensors:*

$$T_{\alpha j}^{m} = -G_{j\alpha}^{m}, \quad P_{i(j)}^{m(\beta)} = C_{i(j)}^{m(\beta)} = 0, \quad P_{(\mu)i(j)}^{(m)\ (\beta)} = \frac{\partial N_{(\mu)i}^{(m)}}{\partial x_{\beta}^{j}} - \delta_{\mu}^{\beta} L_{ij}^{m} = 0,$$

$$P_{(\mu)\alpha(j)}^{(m)\ (\beta)} = \frac{\partial M_{(\mu)\alpha}^{(m)}}{\partial x_{\beta}^{j}} - \delta_{\mu}^{\beta} G_{j\alpha}^{m} + \delta_{j}^{m} H_{\mu\alpha}^{\beta} = -\delta_{\mu}^{\beta} G_{j\alpha}^{m},$$

$$R_{(\mu)\alpha\beta}^{(m)} = \frac{\delta M_{(\mu)\alpha}^{(m)}}{\delta t^{\beta}} - \frac{\delta M_{(\mu)\beta}^{(m)}}{\delta t^{\alpha}}, \quad R_{(\mu)\alpha j}^{(m)} = \frac{\delta M_{(\mu)\alpha}^{(m)}}{\delta x^{j}} - \frac{\delta N_{(\mu)j}^{(m)}}{\delta t^{\alpha}},$$

$$R_{(\mu)ij}^{(m)} = \frac{\delta N_{(\mu)i}^{(m)}}{\delta x^{j}} - \frac{\delta N_{(\mu)j}^{(m)}}{\delta x^{i}}, \quad S_{(\mu)(i)(j)}^{(m)(\alpha)(\beta)} = \delta_{\mu}^{\alpha} C_{i(j)}^{m(\beta)} - \delta_{\mu}^{\beta} C_{j(i)}^{m(\alpha)} = 0.$$

Using the general formulas from the book [6], which give the curvature d-tensors of a generalized multi-time Lagrange space, we obtain the following result:

Theorem 6 *The generalized multi-time Lagrange space $\mathcal{QLG}ML_p^n$ is characterized by the following adapted curvature d-tensors:*

$$H^{\alpha}_{\eta\beta\gamma} = \frac{\partial H^{\alpha}_{\eta\beta}}{\partial t^{\gamma}} - \frac{\partial H^{\alpha}_{\eta\gamma}}{\partial t^{\beta}} + H^{\mu}_{\eta\beta}H^{\alpha}_{\mu\gamma} - H^{\mu}_{\eta\gamma}H^{\alpha}_{\mu\beta},$$

$$R^{l}_{i\beta\gamma} = \frac{\partial G^{l}_{i\beta}}{\partial t^{\gamma}} - \frac{\partial G^{l}_{i\gamma}}{\partial t^{\beta}} + G^{m}_{i\beta}G^{l}_{m\gamma} - G^{m}_{i\gamma}G^{l}_{m\beta},$$

$$R^{l}_{i\beta k} = \frac{\partial G^{l}_{i\beta}}{\partial x^{k}} - \frac{\partial \Gamma^{l}_{ik}}{\partial t^{\beta}} + G^{m}_{i\beta}\Gamma^{l}_{mk} - \Gamma^{m}_{ik}G^{l}_{m\beta},$$

$$R^{l}_{ijk} = \frac{\partial \Gamma^{l}_{ij}}{\partial x^{k}} - \frac{\partial \Gamma^{l}_{ik}}{\partial x^{j}} + \Gamma^{m}_{ij}\Gamma^{l}_{mk} - \Gamma^{m}_{ik}\Gamma^{l}_{mj},$$

$$P^{l\ (\gamma)}_{i\beta(k)} = 0, \quad P^{l\ (\gamma)}_{ij(k)} = 0, \quad S^{l(\beta)(\gamma)}_{i(j)(k)} = 0.$$

4 Generalized multi-time field theories on the space $\mathcal{QLG}ML_p^n$

4.1 Multi-time gravitational field

The fundamental vertical metrical d-tensor of the space $\mathcal{QLG}ML_p^n$ naturally induces the multi-time gravitational h-potential G, defined by

$$G = h_{\alpha\beta}dt^{\alpha} \otimes dt^{\beta} + g_{ij}dx^{i} \otimes dx^{j} + h^{\alpha\beta}g_{ij}\delta x^{i}_{\alpha} \otimes \delta x^{j}_{\beta}.$$

We postulate that the generalized Einstein equations, that govern the multi-time gravitational h-potential of the space $\mathcal{QLG}ML_p^n$, are the equations

$$Ric(C\Gamma) - \frac{Sc(C\Gamma)}{2}G = \mathcal{K}\mathcal{T}, \tag{3}$$

where $Ric(C\Gamma)$ represents the Ricci d-tensor associated with the generalized Cartan connection, $Sc(C\Gamma)$ is the scalar curvature, $\mathcal{K}$ is the Einstein constant and $\mathcal{T}$ is the stress-energy d-tensor of matter.

Using now the general formulas from the book [6], we find the following geometrical results:

Proposition 7 *The Ricci d-tensor $Ric(C\Gamma)$ of the space $\mathcal{QLG}ML_p^n$ is determined by the following adapted components:*

$$R_{\alpha\beta} := H_{\alpha\beta} = H^{\mu}_{\alpha\beta\mu}, \quad R^{(\alpha)}_{i(j)} := P^{(\alpha)}_{i(j)} = -P^{m\ (\alpha)}_{im(j)} = 0,$$

$$R^{(\alpha)}_{(i)j} := P^{(\alpha)}_{(i)j} = P^{m\ (\alpha)}_{ij(m)} = 0, \quad R^{(\alpha)}_{(i)\beta} := P^{(\alpha)}_{(i)\beta} = P^{m\ (\alpha)}_{i\beta(m)} = 0,$$

$$R^{(\alpha)(\beta)}_{(i)(j)} := S^{(\alpha)(\beta)}_{(i)(j)} = S^{m(\beta)(\alpha)}_{i(j)(m)} = 0, \quad R_{i\alpha} = R^{m}_{i\alpha m}, \quad R_{ij} = R^{m}_{ijm}.$$

Corollary 8 *The scalar curvature $Sc(C\Gamma)$ of the space $\mathcal{QLGML}_p^n$ is given by*

$$Sc(C\Gamma) = H + R,$$

where $H = h^{\alpha\beta}H_{\alpha\beta}$ and $R = g^{ij}R_{ij}$.

Consequently, the main result of the generalized Riemann-Lagrange geometry of the multi-time gravitational field is

Theorem 9 *The global generalized Einstein equations (3), that govern the space $\mathcal{QLGML}_p^n$, have the local form*

$$\begin{cases} H_{\alpha\beta} - \dfrac{H+R}{2}h_{\alpha\beta} = \mathcal{KT}_{\alpha\beta} \\ R_{ij} - \dfrac{H+R}{2}g_{ij} = \mathcal{KT}_{ij} \\ -\dfrac{H+R}{2}h^{\alpha\beta}g_{ij} = \mathcal{KT}^{(\alpha)(\beta)}_{(i)(j)} \\ 0 = \mathcal{T}_{\alpha i}, \qquad R_{i\alpha} = \mathcal{KT}_{i\alpha}, \quad 0 = \mathcal{T}^{(\alpha)}_{(i)\beta} \\ 0 = \mathcal{T}_{\alpha(i)}^{\ (\beta)}, \quad 0 = \mathcal{T}^{(\alpha)}_{i(j)}, \qquad 0 = \mathcal{T}^{(\alpha)}_{(i)j}, \end{cases}$$

where $\mathcal{T}_{AB}$, $A, B \in \left\{\alpha,\ i,\ {}^{(\alpha)}_{(i)}\right\}$ are the adapted components of the stress-energy d-tensor $\mathcal{T}$.

4.2 The multi-time electromagnetism

In order to expose the multi-time electromagnetic theory of the space $\mathcal{QLGML}_p^n$, we construct the *metrical deflection d-tensors*

$$D^{(\alpha)}_{(i)j} = \left[\mathcal{G}^{(\alpha)(\mu)}_{(i)(m)}x^m_\mu\right]_{|j} = -\frac{h^{\alpha\mu}}{2}\frac{\partial g_{ij}}{\partial t^\mu},$$
$$d^{(\alpha)(\beta)}_{(i)(j)} = \left[\mathcal{G}^{(\alpha)(\mu)}_{(i)(m)}x^m_\mu\right]|^{(\beta)}_{(j)} = h^{\alpha\beta}g_{ij}.$$

In such a context, we can introduce the *electromagnetic 2-form* of the space $\mathcal{QLGML}_p^n$, by setting

$$F = F^{(\alpha)}_{(i)j}\delta x^i_\alpha \wedge dx^j + f^{(\alpha)(\beta)}_{(i)(j)}\delta x^i_\alpha \wedge \delta x^j_\beta,$$

where

$$F^{(\alpha)}_{(i)j} = \frac{1}{2}\left[D^{(\alpha)}_{(i)j} - D^{(\alpha)}_{(j)i}\right] = 0, \qquad f^{(\alpha)(\beta)}_{(i)(j)} = \frac{1}{2}\left[d^{(\alpha)(\beta)}_{(i)(j)} - d^{(\alpha)(\beta)}_{(j)(i)}\right] = 0.$$

Because $F = 0$, we deduce that the multi-time electromagnetic theory of the space $\mathcal{QLGML}_p^n$ is trivial.

References

[1] G.S. Asanov, *Jet extension of Finslerian gauge approach*, Fortschritte der Physik, vol. **38**, no. **8** (1990), 571-610.

[2] G. Giachetta, L. Mangiarotti, G. Sardanashvily, *Covariant Hamitonian field theory,* arXiv:hep-th/9904062, (1999).

[3] M. Gotay, J. Isenberg, J.E. Marsden, R. Montgomery, *Momentum maps and classical relativistic fields. Part I: Covariant field theory*, arXiv:physics/9801019v2 [math-ph], (2004).

[4] D.D. Holm, J.E. Marsden, T.S. Raţiu, *The Euler-Poincaré equations and semidirect products with applications to continuum theories,* arXiv:chao-dyn/9801015, (1998).

[5] R. Miron, M.S. Kirkovits, M. Anastasiei**,** *A geometrical model for variational problems of multiple integrals*, Proc. of Conf. on Diff. Geom. and Appl., Dubrovnik, Yugoslavia, June 26-July 3 (1988), 8-25.

[6] M. Neagu, *Riemann-Lagrange Geometry on* 1*-Jet Spaces*, Matrix Rom, Bucharest, 2005.

[7] P.J. Olver, *Canonical elastic moduli*, Journal of Elasticity **19** (1988), 189-212.

Mircea Neagu
Transilvania University of Braşov
Department of Mathematics and Informatics
Blvd. Iuliu Maniu 50, 500091 Braşov, Romania.
E-mail: mircea.neagu@unitbv.ro